DC Instalift[™]Goji (glycopeptides)

INCI: Water (and) Butylene Glycol (and) Lycium Barbarum Fruit Extract **

March 5, 2013 rev.

Immediate wrinkle lifting and firming / Long-term anti-aging benefits

Lycium Barbarum, also referred to Tibetan Goji berry or Wolfberry, is a member of the nightshade family of plants. This species of plant is a deciduous woody perennial, native to southeastern Europe and Asia.

Goji has long played an important role in traditional Chinese medicine where it is known to enhance immune system function, improve eyesight, protect the liver and improve circulation, among other effects. In recent years, it has become highly prized in the global health food market with attention being focused on their nutrient value and antioxidant content (carotenoids (beta-carotene and zeaxanthin), polysaccharides, vitamins, minerals and essential fatty acids).



DC Instalift Goji is an isolated active glycopeptide fraction from the Goji berry which combines the skin lifting and firming strength of both proteins and polysaccharides. This specific glycopeptide fraction forms a perfect rigid lifting mesh to support sagging and wrinkled skin for creating an immediate visible lifting appearance. DC Instalift™ Goji further offers critical long term anti-aging functions via: inhibition of MMP 1 expression, antioxidant efficacy, anti-apoptotic activity, and collagen stimulation for a complete anti-aging treatment.

BENEFITS	*	Smoothing	•	Firming	
	•	Lifting	•	Free Radical Scavenging	
	•	Moisturizing	•	Anti-Aging	
		lifting mid	cro ripples	••	
APPLICATIONS	*	Forehead wrinkles	•	Frown lines	
	•	Reduction of crow's feet	•	Sagging skin areas	
	•	Eye brow lift	•	Body firming products	
TYPICAL PROPERTIES					
Appearance		Dark Brown Liquid			
рН		5.00 - 7.00			
Solids		7.00-8.00% Lycium Barbarum Fruit	Extract		
FORMULATION GUIDELINES					-
Recommended use level		3%			
pH range		4-8			
** Glycol-free version available: DC	Instalift [™] Goj	i GF			




DC Instalift[™]Goji (glycopeptides)

INCI: Water (and) Butylene Glycol (and) Lycium Barbarum Fruit Extract **

Page 2 of 2

TIGHTENING EFFECT

DC Instalift $^{\rm TM}$ Goji forms tightening micro-ripples observed after 1 hour on slide at 400X






DC Instalift™Goji at T=0, 400x magnification, sample is left of meniscus line

DC Instalift™Goji at T=1hr, 400x magnification, sample is left of meniscus line

ANTI-AGING EFFECT

Inhibition of MMP-1 Expression by increasing concentrations of Goji Glycopeptides




■ 0% Goji Glycopeptides ■ 0.01% Goji Glycopeptides □ 0.10% Goji Gycopeptides □ 1% Goji Glycopeptides

DC Instalift $^{\text{TM}}$ Goji (goji glycopeptides) further offers critical long term anti-aging functions via: inhibition of MMP 1 expression, antioxidant activity, anti-apoptotic, and collagen stimulatory activity for a complete anti-aging treatment.

ANTIOXIDANT EFFECT


This graph demonstrates the protective antioxidant properties of goji glycopeptides and its capability of inhibiting the oxidative capacity of superoxide free radical demonstrated by measuring the rate of cytochrome C reduction.



At increasing concentrations of goji glycopeptides the rate of oxidation is significantly diminished.


SMOOTHING EFFECT

Baseline vs. 60 minutes after application of 3% DC Instalift™ Goji in water



FIRMING EFFECT

DC Instalift™Goji was tested at 3% in water on 19 subjects using Twistometery via Dermal Torque Meter Method. Results are expressed as % variation after one hour compared to a baseline measurement.



Placebo (water) showed no significant change after 1 hour. DC Instalift™ Goji improved skin firmness by 26.8% after 1 hour with statistical significance p<0.001

REFERENCES

- Protective effect of Fructus Lycii polysaccharides against time and hyperthermia-induced damage in cultured seminiferous epithelium; Yanrong Wang, Hui Zhao, Xinsheng Sheng, Patricia E. Gambino, Brian Costello, Krzysztof Bojanowski; Journal of Ethnopharmacology 82 (2002) 169/175
- Lycium barbarum glycoconjugates: effect on human skin and cultured dermal fibroblasts, H. Zhao, A. Alexeev, E. Chang, G. Greenburg, K. Bojanowski; Phytomedicine 12 (2005) 131–137

The information contained in this technical bulletin is presented in good faith, and to the best of our knowledge believed to be true and accurate. No representations or warranties, expressed or implied is made or intended. Information is supplied upon the condition that the persons receiving same will make their own determination as to its suitability for their purposes prior to use. No recommendation should be construed as an inducement to use a material in infringement of patents or applicable government regulations. In no event will Resources of Nature be responsible or liable for any loss of profits, lost goodwill, direct, special, indirect, incidental, or consequential damages of any nature whatsoever.





DISTRIBUTED BY

UNIT 2 CORINIUM INDUSTRIAL ESTATE

RAANS ROAD, AMERSHAM

HP6 6JQ, UNITED KINGDOM

TEL: +44 (0) 1494 410160
FAX: +44 (0) 1494 722320
EMAIL: SALES®AMMEREX.COM